

Figure-1a: chromatogram *Ficus capensis* leaves extract (MWE) 280 nm.

1: gallic acid, 2: protocatechuic acid, 3: catechin, 4: caffeic acid, 5: epicatechin, 6: *p*-coumaric acid, 7: rutin, 10: cinnamic acid, 11: quercetin




Figure-1b: chromatogram *Ficus capensis* leaves extract (MWE) 320 nm.

1: gallic acid, 2: protocatechuic acid, 4: caffeic acid, 5: epicatechin, 6: *p*-coumaric acid, 8: quercetin glucosyl, 10: cinnamic acid, 11: quercetin




Figure-2a: chromatogram WE F. capensis leaves analysis (280nm).

5: epicatechin, 6: p - coumaric acid, 8: quercetin glucosyl, 9: quercetin dihydrat, 10: cinnamic acid, 11: quercetin




Figure-2b: WE chromatogram F. capensis (320 nm).

2: protocatechiuc acid, 3: catechin, 5: epicatechin, 6: p-coumaric acid, 7: rutin, 8: quercetin glucosyl, 9: quercetin dihydrat, 10: cinnamic acid




Figure-3: Relationship between inhibition percentage of NO_2^- and the concentration (volume) of $[\text{NO}_2]^-$.
 IP: % Inhibition